

Pulvex
Tick & Flea
SPOT

Reg. No. G3923 Act 36 of 1947

FOR EXTERNAL ANIMAL USE ONLY

for puppies

Pulvex®

DO NOT USE ON CATS
STORE BELOW 25°C.

WARNINGS

Handle with care. Avoid contact with the eyes. Store away from food and feed. Keep out of reach of children and uninformed persons. Dangerous for fish so do not contaminate ponds, dams or rivers with the product or packaging. Dispose of packaging safely in household refuse bin. Do not allow treated animals to swim for 12 hours after treatment. Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

Avoid contamination of food, drinking water and food utensils. Whilst handling Pulvex® Tick & Flea Spot, avoid the solution coming into contact with the fingers. Do not handle the area of application for 3-6 hours after treatment. Wash hands and any splashes immediately after use. If treating a number of dogs, protect your hands by wearing gloves. Should the dog experience any minor irritation on the skin at the site of application, bathe the dog with a mild non-insecticidal shampoo. Cats must not groom the dog where the solution has been applied. If this occurs, separate the animals and consult your vet. It can be beneficial to apply the product in the evening, but do not allow the dog to sleep with people, especially children, within 3-6 hours after treatment.

DIRECTIONS FOR USE
USE ONLY AS DIRECTED.

Always keep the sachet with the box, until used.
Open one sachet and remove pipette. With one hand, hold the pipette in an upright position away from your face. Snap tip open by bending and folding it back on itself. Apply by parting the dog's coat on the back between the shoulder blades and apply directly onto the skin and to the tips of the ears if flies are a problem. Avoid applying the solution to the hair and do not rub into the skin as this will reduce the product's efficacy. For dogs and puppies weighing up to and including 15kg (i.e. Cocker Spaniel), apply 1 x 1ml pipette onto the skin between the shoulders and the ears, if needed. For dogs and puppies over 15kg, apply 2 pipettes. Apply one onto the skin between the shoulder blades, and ears if needed for fly control. Apply the other pipette onto the skin on the back at the base of the tail. **Pulvex® Tick & Flea Spot** should not be used on puppies less than 2 weeks old and should not be used on cats, as this product is TOXIC TO CATS.

FREQUENCY OF USE

Pulvex® Tick & Flea Spot for dogs will provide more than 4 weeks of control and protection against flies, ticks (kennel ticks/brown dog ticks) and fleas. In the case of severe infestations, it is safe to use a further treatment 7 days after the initial dose. If the treated dog gets very wet from shampooing or swimming, the protection will be reduced and the solution should be re-applied, but not before 7 days.

See bottom of box for Batch No./Expiry Date

NOT FOR USE ON CATS OR PUPPIES
UNDER TWO WEEKS OLD

FOR EXTERNAL ANIMAL USE ONLY

Pulvex®

Tick & Flea
SPOT

Reg. No. G3923 Act 36 of 1947

for puppies
from 2 weeks old

kills and controls flies, ticks
(kennel ticks/brown dog ticks)
and fleas for more than 4 weeks

DO NOT USE ON CATS

Contains: Permethrin 65% m/v

kills
& controls



®Registered Trademark of
Intervet (Pty) Ltd.

Registration Holder: Model
Product Distributors (Pty) Ltd.
Co. Reg No. 1997/005938/07
PO Box 293,
Modderfontein, 1645

Tel: 011 997 3400 or
08600 MODEL (66335)

1x 1ml Pipette



Line: 002203, Tick & Flea Spot, PUPPES, 1/13

BATCH NO:

EXPIRY DATE: